

Intended for healthcare professionals

Account Administrators: Review your remote access options for SAGE Journals


Journal of Psychopharmacology

Associations between lifetime classic psychedelic use and markers of physical health

Otto Simonsson , James D Sexton, Peter S Hendricks

First Published March 9, 2021 | Research Article | [Find in PubMed](#)

<https://doi.org/10.1177/0269881121996863>



Abstract

Background:

In recent years, there has been significant research on the mental health effects of classic psychedelic use, but there is very little evidence on how classic psychedelics might influence physical health.

Aims:

The purpose of the present study was to investigate the associations between lifetime classic psychedelic use and markers of physical health.

Methods:

Using data from the National Survey on Drug Use and Health (2015-2018) with 171,766 (unweighted) adults aged 18 or above in the United States, the current study examined the associations between lifetime classic psychedelic use and three markers of physical

health (self-reported overall health, body mass index, and heart condition and/or cancer in the past 12 months) while controlling for a range of covariates.

Results:

Respondents who reported having tried a classic psychedelic at least once in their lifetime had significantly higher odds of greater self-reported overall health and significantly lower odds of being overweight or obese versus having a normal weight. The association between lifetime classic psychedelic use and having a heart condition and/or cancer in the past 12 months approached conventional levels of significance, with lower odds of having a heart condition and/or cancer in the past 12 months for respondents who had tried a classic psychedelic at least once.

Conclusion:

The results of the present study suggest that classic psychedelics may be beneficial to physical health. Future research should investigate the causal effects of classic psychedelics on physical health and evaluate possible mechanisms.

Keywords

[Classic psychedelics](#), [psilocybin](#), [LSD](#), [health](#), [body mass index](#), [cancer](#), [heart disease](#)

Introduction

The effects of psychedelic drug use on human cognition and behaviour have recently received significant scientific attention ([Rucker et al., 2018](#); [Sessa, 2018](#)). The research has been conducted in controlled settings and has primarily focused on classic psychedelics, which are a subclass of psychedelics, with little evidence of physiological toxicity, known to act as agonists primarily at 5-HT_{2A} receptors ([dos Santos et al., 2018](#); [Johnson et al., 2018](#); [Winkelman, 2014](#)). The three main classes of classic psychedelics (tryptamines, lysergamides and phenethylamines) are distinguished by unique chemical structures and neurochemical mechanisms ([Szabo, 2015](#)). Most notably, they include N,N-dimethyltryptamine (DMT), the DMT-containing admixture ayahuasca, psilocybin, lysergic acid diethylamide (LSD), mescaline, and the mescaline-containing cacti peyote and San Pedro ([Sexton et al., 2019a](#)).

The evidence to date suggests that classic psychedelics have immunomodulatory and anti-inflammatory properties ([Flanagan and Nichols, 2018](#); [Freckska et al., 2016](#); [Nichols,](#)

2009; Szabo, 2015, 2019); carry low risk of adverse effects when administered by health professionals in a safe and supportive environment (Nutt and Carhart-Harris, 2020; Nutt et al., 2010; Rucker et al., 2018); and can be effective in the treatment of depression, anxiety and addiction (Aday et al., 2020; Carhart-Harris et al., 2016, 2018; Davis et al., 2020; Goldberg et al., 2020; Johnson et al., 2014; Krebs and Johansen, 2012; Luoma et al., 2020). For example, patients with treatment-resistant depression experienced reductions in depressive symptoms after two oral doses of psilocybin. There was no control group in the study, but the depressive symptoms remained significantly reduced at 1 week, 3 months, and 6 months post-treatment (Carhart-Harris et al., 2016, 2018). The patients were interviewed post-treatment and many of them reported significant improvements in health behaviour (Watts et al., 2017), which suggests that classic psychedelic use might induce behavioural changes favourable to physical health.

While double-blind, randomized, placebo-controlled trials are warranted to experimentally examine the effects of classic psychedelics on physical health, population studies can provide insight into these knowledge gaps. The National Survey on Drug Use and Health (NSDUH) has frequently been analyzed to provide weighted estimates of the prevalence and associations of lifetime classic psychedelic use in the United States. The findings have varied from 13.4 to 13.6% of the adult population reporting lifetime classic psychedelic use (Hendricks et al., 2017; Johansen and Krebs, 2015) and have consistently showed that lifetime classic psychedelic users are more likely to be male, white, younger than 65 years of age, and have higher income and education (Johansen and Krebs, 2015; Krebs and Johansen, 2013a, 2013b). The results have also shown that lifetime classic psychedelic use is associated with lower odds of psychological distress and suicidality (Hendricks et al., 2015), lower odds of opioid abuse and dependence (Pisano et al., 2017), and lower odds of criminal behavior (Hendricks et al., 2017), which broadly mirrors the research that suggests therapeutic efficacy of classic psychedelics as well as the low risk of harm to self and others that classic psychedelics have been ascribed by drug experts in the United Kingdom (Nutt et al., 2010), the Netherlands (Van Amsterdam et al., 2010) and Australia (Bonomo et al., 2019). There have, in other words, been several population studies on lifetime classic psychedelic use, but the association between lifetime classic psychedelic use and physical health remains unexplored.

Using pooled data from the NSDUH (2015–2018), the present study seeks to investigate the association between lifetime classic psychedelic use and three markers of physical health (self-reported overall health, body mass index (BMI), and heart condition and/or cancer in past 12 months). We hypothesized that lifetime classic psychedelic use would be associated with better physical health status.

Materials and methods

Data and population

The NSDUH is an annual survey designed to measure the prevalence of substance use and mental health issues in the United States. The present study used pooled data from the NSDUH survey years 2015 to 2018, which were weighted to reflect the civilian noninstitutionalized population and contained responses from 171,766 (unweighted) adults aged 18 or above. The NSDUH sampling and questionnaire methodology are described on their website: https://nsduhweb.rti.org/respweb/about_nsduh.html

Variables

The dependent variables were self-reported overall health (variable HEALTH2 recoded; 1 = Fair/Poor, 2 = Good, 3 = Very Good, 4 = Excellent), BMI (variable BMI2 recoded per National Institute of Health (NIH) guidelines ([National Heart, Lung, and Blood Institute \(NHLBI\) 1998](#)): 1 = Underweight (<18.5), 2 = Normal Weight (18.5–25), 3 = Overweight (25–30), 4 = Obesity – Class 1 (30–35), 5 = Obesity – Class 2 (35–40), 6 = Extreme Obesity – Class 3 (>40)), and heart condition and/or cancer in past 12 months (variables HRTCONDYR and CANCERYR combined such that a ‘yes’ response to either variable was coded as 1 = Yes whereas a ‘no’ response to both variables was coded as 0 = No).

The independent variables were DMT (code 616 from variables HALLUCOT1, HALLUCOT2, HALLUCOT3, HALLUCOT4 and HALLUCOT5), ayahuasca (an entheogenic brew that contains DMT; code 6103 from variables HALLUCOT1, HALLUCOT2, HALLUCOT3, HALLUCOT4 and HALLUCOT5), psilocybin (variable PSILCY2 = 1), LSD (variable lsdflag = 1), mescaline (variable MESC2 = 1), and peyote or San Pedro (cacti that contain mescaline; variable PEYOTE2 = 1 or code 6077 from variables HALLUCOT1, HALLUCOT2, HALLUCOT3, HALLUCOT4 and HALLUCOT5). Respondents reporting that they had ever, even once, used any of the above classic psychedelics were coded as positive for lifetime classic psychedelic use, whereas those indicating that they had never used any of these substances were coded as negative. The question in the NSDUH concerning use of DMT, alpha-methyltryptamine (AMT) and 5-methoxy-N, N-diisopropyltryptamine (5-MeO-DIPT) (variable DAMTFXFLAG: ‘Have you ever, even once, used any of the following: DMT, also called dimethyltryptamine, AMT, also called alpha-methyltryptamine, or Foxy, also called 5-MeO-DIPT?’) was not included, because neither AMT nor 5-MeO-DIPT are classified as classic psychedelic and DMT use alone could not be determined from the question.

The control variables were age in years (variable CATAG6; 18–25, 26–34, 35–49, 50–64 or 65 or older); sex (variable IRSEX; male or female); sexual orientation (variable SEXIDENT; heterosexual, lesbian or gay, bisexual); ethnoracial identity (variable NEWRACE2; non-Hispanic White, non-Hispanic African American, non-Hispanic Native American/Alaska Native, non-Hispanic Native Hawaiian/Pacific Islander, non-Hispanic Asian, non-Hispanic more than one race or Hispanic); educational attainment (variable IREDUHIGHST2; 5th grade or less, 6th grade, 7th grade, 8th grade, 9th grade, 10th grade, 11th or 12th grade completed, High school diploma/GED, some college credit but no degree, Associate's degree, College degree or higher), annual household income (variable INCOME; less than US\$20,000, US\$20,000–49,999, US\$50,000–74,999 or US\$75,000 or more); marital status (variables IRMARITSTAT and IRMARIT recoded; married, divorced/separated, widowed or never married); self-reported engagement in risky behaviour (variable RSKYFQTES recoded; never, seldom, sometimes or always), lifetime cocaine use (variable COCFLAG; ever used or never used), lifetime other stimulant use (variable STMANYFLAG; ever used or never used), lifetime sedative use (variable SEDANYFLAG; ever used or never used), lifetime tranquilizer use (variable TRQANYFLAG; ever used or never used), lifetime heroin use (variable HERFLAG; ever used or never used), lifetime pain reliever use (variable PNRANYFLAG; ever used or never used), lifetime marijuana use (variable MRJFLAG; ever used or never used), lifetime phencyclidine (PCP) use (variable PCPFLAG; ever used or never used), lifetime 3,4-methylenedioxymethamphetamine (MDMA/ecstasy) use (variable ECSTMOFLAG; ever used or never used), lifetime inhalant use (variable INHALFLAG; ever used or never used), lifetime smokeless tobacco use (variable SMKLSFLAG; ever used or never used), lifetime pipe tobacco use (variable PIPFLAG; ever used or never used), lifetime cigar use (variable CGRFLAG; ever used or never used), lifetime daily cigarette use (variable CDUFLAG; ever used or never used) and age of first alcohol use (variable IRALCAGE recoded; less than 13 years of age (Preteen), 13–19 years of age (Teen), more than 19 years of age (Adult), or never used). These control variables were coded as separate covariates and broadly mirror the covariates of prior investigations ([Hendricks et al., 2015](#); [Sexton et al., 2019b](#)), except for lifetime smokeless tobacco, pipe tobacco, cigar and daily cigarette use, as well as age of first alcohol use, which were added to control for a lifetime history of major health risk factors ([Christensen et al., 2018](#); [Hu et al., 2017](#); [Inoue-Choi et al., 2019a, 2019b](#); [Levola et al., 2020](#)). Lastly, a recoded version of the Kessler Psychological Distress Scale ([Kessler et al. 2002, 2010](#)) was also included as a control variable (variable K6SCMON recoded into dichotomous variable), but only in the ordered logistic regression model predicting self-reported overall health (see below) to ensure that

self-reported overall health was not influenced by the mental health status of the respondents.

Statistical analyses

The present study used descriptive statistics to present an overview of lifetime use of classic psychedelics in the United States (Table 1). Multiple regressions were used to calculate adjusted odds ratios with 95% confidence intervals and examine the unique associations between lifetime classic psychedelic use and markers of physical health. Ordered logistic regression was used to examine the association between lifetime classic psychedelic use and self-reported overall health (Table 2); multinomial logistic regression was used to examine the association between lifetime classic psychedelic use and BMI (Table 3); and logistic regression was used to examine the association between lifetime classic psychedelic use and having a heart condition and/or cancer in the past 12 months (Table 4).



Table 1. Lifetime classic psychedelic (DMT, Ayahuasca, Psilocybin, LSD, Mescaline, Peyote or San Pedro) users in the United States (2015–2018).



Table 2. Lifetime classic psychedelic use and self-reported overall health.




Table 3. Lifetime classic psychedelic use and body mass index.




Table 4. Lifetime classic psychedelic use and heart condition and/or cancer in the past year.

The analyses used weights provided by the NSDUH, and the control variables listed above were included as covariates in the regression models to control for potential sources of confounding. Insofar that the NSDUH is a nationally representative survey, there was no a priori rationale for identifying or removing outliers. The NSDUH conducts statistical imputation and revision for missing values for select variables, denoted by ‘IMP’

or 'IR' prefixes. A number of constraints are put in place by the NSDUH to ensure consistency in imputed values with non-missing values for use in multivariate analyses. In the present analyses we did not conduct additional imputations beyond what the NSDUH has already provided in their annual releases. All missing values were treated as missing. Additional information on the NSDUH imputation procedure can be found in the 'Statistical Imputation' section in the introduction of each annual codebook. Finally, though there was no control for multiple comparisons, exact p-values are reported to the fourth decimal place, which allows for the application of conservative Bonferroni-type corrections of the reader's choosing ([SAMHSA, 2019](#)).

Results

Frequency distributions

[Table 1](#) presents descriptive statistics of lifetime classic psychedelic users in the United States (2015–2018). As shown in the table, approximately 14% of the sample reported lifetime classic psychedelic use, which suggests that almost 34 million American adults have used a classic psychedelic at least once in their lifetime, based on the population estimates from the NSDUH.

Multiple regressions

[Table 2](#) presents results from the ordered logistic regression on the association between lifetime classic psychedelic use and self-reported overall health. As illustrated in the table, lifetime classic psychedelic use was associated with significantly higher odds of greater self-reported overall health.

[Table 3](#) presents results from the multinomial logistic regression on the association between lifetime classic psychedelic use and BMI. As illustrated in the table, lifetime classic psychedelic use was associated with significantly lower odds of being overweight or obese as compared to having a normal weight.

[Table 4](#) presents results from the logistic regression on the association between lifetime classic psychedelic use and having a heart condition and/or cancer in the past year. As illustrated in the table, the association between lifetime classic psychedelic use and having a heart condition and/or cancer in the past 12 months approached conventional levels of significance, with lower odds of having a heart condition and/or cancer in the past 12 months for respondents who had tried a classic psychedelic at least once.

Discussion

The present study investigated the association between lifetime classic psychedelic use and three markers of physical health (self-reported overall health, BMI, and heart condition and/or cancer in the past 12 months). Findings show that respondents who reported having ever used a classic psychedelic had significantly higher odds of greater self-reported overall health and significantly lower odds of being overweight or obese as compared to having a normal weight. The association between lifetime classic psychedelic use and having a heart condition and/or cancer in the past 12 months approached conventional levels of significance, with lower odds of having a heart condition and/or cancer in the past 12 months for respondents who had tried a classic psychedelic at least once. Taken together, these results suggest that classic psychedelics may have long-term beneficial effects beyond improved mental health.

While the acute transcendent experience occasioned by classic psychedelics may presumably induce long-term changes in health behaviour that contribute to better physical health, it is plausible that there are other key mechanisms through which classic psychedelics could influence physical health, including improvements on various indices of mental health beyond the simple absence of psychological distress (e.g. increased prosociality, trait mindfulness and purpose in life; [Griffiths et al., 2018](#); [Murphy-Beiner and Soar, 2020](#)), many of which are well-known risk factors for physical maladies ([Chaddha et al., 2016](#); [Germann, 2020](#); [Hernandez et al., 2018](#)); immunomodulatory and anti-inflammatory effects of relevance to physical health ([Flanagan and Nichols, 2018](#); [Frecka et al., 2013, 2016](#); [Szabo, 2015, 2019](#); [Szabo et al., 2014](#); [Thompson and Szabo, 2020](#); [Tourino et al., 2013](#); [Winkelman and Sessa, 2019](#)); and high affinity to receptor subtypes (e.g. serotonin 2A receptors) that are implicated in the pathophysiology of different physical disorders ([Nichols, 2009](#); [Thompson and Szabo, 2020](#)). Future research is needed to better understand potential causal pathways of classic psychedelics on physical health.

There are several limitations with the present study that need serious consideration before the results are interpreted. First, the cross-sectional design of the study limits causal inference. The analyses controlled for multiple sources of potential confounding, but the associations might have been obscured by response bias or latent variables that were not controlled for (e.g. a common factor predisposing one to classic psychedelic use may also predispose one to healthy lifestyle behaviours including physical activity). Second, the dataset did not contain information on frequency of classic psychedelic use, dose used or context of use. The present study could therefore not evaluate frequency, dose or context-


specific relationships between classic psychedelic use and physical health markers. Third, it is also not possible to rule out that classic psychedelic use might have caused harm on the individual level, even if it did not obfuscate the population-level associations. Fourth, given the potential importance of immunomodulatory and inflammatory factors in the current study, it would have been sensible to also control for regular anti-inflammatory drug (e.g. nonsteroidal anti-inflammatory drug (NSAID)) use, but assessment of this behaviour was not included in the NSDUH. Fifth, BMI has been widely used as a screening tool for overweight or obesity, but it does not account for details such as fat distribution, which limits its utility as a marker of physical health (Prentice and Jebb, 2001). Finally, it is noted that some associations of lifetime classic psychedelic use were somewhat modest in size (e.g. heart condition and/or cancer in the past year). However, even modest effects can have substantial impacts at the population level. For instance, considering approximately 1.2 million people die from heart disease or cancer every year in the United States alone (Heron, 2019), even a small decrease (e.g. 11%) in the prevalence of these illnesses could translate to thousands of lives saved annually.

Conclusion

The psychedelic research to date has primarily focused on mental health, but relatively little is known about how classic psychedelics might influence physical health. The findings in the present study suggest that lifetime classic psychedelic use is associated with higher odds of better physical health status, which demonstrates the need for more rigorous research to better understand potential causal pathways of classic psychedelics on physical functioning.


Appendix

Logistic regressions



Click to view table

Table A1. Lifetime classic psychedelic use predicting heart condition in the past year.



Click to view table

Table A2. Lifetime classic psychedelic use predicting cancer in the past year.

Multinomial logistic regression



Table A3. Lifetime classic psychedelic use predicting heart condition only, cancer only, and both heart condition and cancer in the past year.

Authors' note

The current study was a secondary analysis of publicly available data files and was exempt from review by the Research Ethics Committee of the Department of Sociology (DREC) at the University of Oxford.

Author contributions

OS conceived of the study and was the primary author who cleaned data, conducted analyses and drafted the manuscript summarizing the findings. JDS commented on draft manuscripts and contributed meaningful expertise to inform methodology and statistical analyses. PSH supervised and commented on draft manuscripts.

Declaration of conflicting interests

PSH is on the scientific advisory board of Bright Minds Biosciences Ltd., Eleusis Benefit Corporation., and Silo Pharma Inc. JDS is the Bioinformatics Director at Eleusis Benefit Corporation.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Otto Simonsson



<https://orcid.org/0000-0003-4197-7566>

References

Aday, JS, Mitzkovitz, CM, Bloesch, EK, et al. (2020) Long-term effects of psychedelic drugs: A systematic review. *Neurosci Biobehav Rev* 113: 179–189.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Bonomo, Y, Norman, A, Biondo, S, et al. (2019) The Australian drug harms ranking study. *J Psychopharmacol* 33: 759–768.

[Google Scholar](#) | [SAGE Journals](#) | [ISI](#)

Carhart-Harris, RL, Bolstridge, M, Day, CMJ, et al. (2018) Psilocybin with psychological support for

treatment-resistant depression: Six-month follow-up. *Psychopharmacology* 235: 399–408.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Carhart-Harris, RL, Bolstridge, M, Rucker, J, et al. (2016) Psilocybin with psychological support for treatment-resistant depression: An open-label feasibility study. *Lancet Psychiatry* 3: 619–627.

[Google Scholar](#) | [Crossref](#) | [Medline](#) | [ISI](#)

Chaddha, A, Robinson, EA, Kline-Rogers, E, et al. (2016) Mental health and cardiovascular disease. *Am J Med* 129: 1145–1148.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Christensen, CH, Rostron, B, Cosgrove, C, et al. (2018) Association of cigarette, cigar, and pipe use with mortality risk in the US population. *JAMA Intern Med* 178: 469–476.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Davis, AK, Barrett, FS, May, DG, et al. (2020) Effects of psilocybin-assisted therapy on major depressive disorder: A randomized clinical trial. *JAMA Psychiatry*. Epub ahead of print 4 November 2020. DOI: [10.1001/jamapsychiatry.2020.3285](https://doi.org/10.1001/jamapsychiatry.2020.3285).

[Google Scholar](#) | [Crossref](#)

dos Santos, RG, Bouso, JC, Alcázar-Córcoles, MÁ, et al. (2018) Efficacy, tolerability, and safety of serotonergic psychedelics for the management of mood, anxiety, and substance-use disorders: A systematic review of systematic reviews. *Expert Rev Clin Pharmacol* 11(9): 889–902.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Flanagan, TW, Nichols, CD (2018) Psychedelics as anti-inflammatory agents. *Int Rev Psychiatry* 30: 363–375.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Frecska, E, Bokor, P, Winkelman, M (2016) The therapeutic potentials of ayahuasca: Possible effects against various diseases of civilization. *Front Pharmacol* 7: 35.

[Google Scholar](#) | [Crossref](#) | [Medline](#) | [ISI](#)

Frecska, E, Szabo, A, Winkelman, MJ, et al. (2013) A possibly sigma-1 receptor mediated role of dimethyltryptamine in tissue protection, regeneration, and immunity. *J Neural Transm* 120: 1295–1303.

[Google Scholar](#) | [Crossref](#) | [Medline](#) | [ISI](#)

Germann, CB (2020) The Psilocybin-Telomere Hypothesis: An empirically falsifiable prediction concerning the beneficial neuropsychopharmacological effects of psilocybin on genetic aging. *Med Hypoth* 134: 109406.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Goldberg, SB, Pace, BT, Nicholas, CR, et al. (2020) The experimental effects of psilocybin on symptoms of anxiety and depression: A meta-analysis. *Psychiatry Res* 284: 112749.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Griffiths, RR, Johnson, MW, Richards, WA, et al. (2018) Psilocybin-occasioned mystical-type

experience in combination with meditation and other spiritual practices produces enduring positive changes in psychological functioning and in trait measures of prosocial attitudes and behaviors. *J Psychopharmacol* 32(1): 49–69.

[Google Scholar](#) | [SAGE Journals](#) | [ISI](#)

Hendricks, PS, Crawford, MS, Cropsey, KL, et al. (2017) The relationships of classic psychedelic use with criminal behavior in the United States adult population. *J Psychopharmacol* 32: 37–48.

[Google Scholar](#) | [SAGE Journals](#)

Hendricks, PS, Thorne, CB, Clark, CB, et al. (2015) Classic psychedelic use is associated with reduced psychological distress and suicidality in the United States adult population. *J Psychopharmacol* 29: 280–288.

[Google Scholar](#) | [SAGE Journals](#) | [ISI](#)

Hernandez, R, Bassett, SM, Boughton, SW, et al. (2018) Psychological well-being and physical health: Associations, mechanisms, and future directions. *Emotion Rev* 10: 18–29.

[Google Scholar](#) | [SAGE Journals](#) | [ISI](#)

Heron, MP (2019) Deaths: Leading causes for 2017. *Natl Vital Stat Rep* 68: 1–77.

[Google Scholar](#) | [Medline](#)

Hu, H, Eaton, WW, Anthony, JC, et al. (2017) Age of first drunkenness and risks for all-cause mortality: A 27-year follow-up from the epidemiologic catchment area study. *Drug Alcohol Depend* 176: 148–153.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Inoue-Choi, M, McNeel, TS, Hartge, P, et al. (2019a) Non-daily cigarette smokers: Mortality risks in the US. *Am J Prevent Med* 56: 27–37.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Inoue-Choi, M, Shiels, MS, McNeel, TS, et al. (2019b) Contemporary associations of exclusive cigarette, cigar, pipe, and smokeless tobacco use with overall and cause-specific mortality in the United States. *JNCI Cancer Spectrum* 3: pkz036.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Johansen, P-Ø, Krebs, TS (2015) Psychedelics not linked to mental health problems or suicidal behavior: A population study. *J Psychopharmacol* 29: 270–279.

[Google Scholar](#) | [SAGE Journals](#) | [ISI](#)

Johnson, MW, Garcia-Romeu, A, Cosimano, MP, et al. (2014) Pilot study of the 5-HT_{2A}R agonist psilocybin in the treatment of tobacco addiction. *J Psychopharmacol* 28: 983–992.

[Google Scholar](#) | [SAGE Journals](#) | [ISI](#)

Johnson, MW, Griffiths, RR, Hendricks, PS, et al. (2018) The abuse potential of medical psilocybin according to the 8 factors of the Controlled Substances Act. *Neuropharmacology* 142: 143–166.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Kessler, RC, Andrews, G, Colpe, LJ, et al. (2002) Short screening scales to monitor population

prevalences and trends in non-specific psychological distress. *Psychol Med* 32: 959–976.

[Google Scholar](#) | [Crossref](#) | [Medline](#) | [ISI](#)

Kessler, RC, Green, JG, Gruber, MJ, et al. (2010) Screening for serious mental illness in the general population with the K6 screening scale: Results from the WHO World Mental Health (WMH) survey initiative. *Intl J Methods Psychiatric Res* 19(S1): 4–22.

[Google Scholar](#) | [Crossref](#) | [Medline](#) | [ISI](#)

Krebs, TS, Johansen, PØ (2012) Lysergic acid diethylamide (LSD) for alcoholism: meta-analysis of randomized controlled trials. *J Psychopharmacol* 26: 994–1002.

[Google Scholar](#) | [SAGE Journals](#) | [ISI](#)

Krebs, TS, Johansen, PØ (2013a) Psychedelics and mental health: A population study. *PLoS One* 8: e63972.

[Google Scholar](#) | [Crossref](#) | [Medline](#) | [ISI](#)

Krebs, TS, Johansen, PØ (2013b) Over 30 million psychedelic users in the United States. *F1000Research* 2: 98.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Levola, J, Rose, RJ, Mustonen, A, et al. (2020) Association of age at first drink and first alcohol intoxication as predictors of mortality: A birth cohort study. *Eur J Public Health* 30: 1189–1193.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Luoma, JB, Chwyl, C, Bathje, GJ, et al. (2020) A Meta-analysis of placebo-controlled trials of psychedelic-assisted therapy. *J Psychoactive Drugs* 52: 289–299.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Murphy-Beiner, A, Soar, K (2020) Ayahuasca's 'afterglow': Improved mindfulness and cognitive flexibility in ayahuasca drinkers. *Psychopharmacology* 237: 1161–1169.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

National Heart, Lung, and Blood Institute (NHLBI) (1998) *Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults: The Evidence Report*. Washington, DC: Government Printing Office.

[Google Scholar](#)

Nichols, CD (2009) Serotonin 5-HT_{2A} receptor function as a contributing factor to both neuropsychiatric and cardiovascular diseases. *Cardiovasc Psychiatry Neurol* 2009: 475108.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Nutt, D, Carhart-Harris, R (2020) The current status of psychedelics in psychiatry. *JAMA Psychiatry* 78: 121–122.

[Google Scholar](#) | [Crossref](#)

Nutt, DJ, King, LA, Phillips, LD (2010) Drug harms in the UK: A multicriteria decision analysis. *Lancet* 376: 1558–1565.

[Google Scholar](#) | [Crossref](#) | [Medline](#) | [ISI](#)

Pisano, VD, Putnam, NP, Kramer, HM, et al. (2017) The association of psychedelic use and opioid use disorders among illicit users in the United States. *J Psychopharmacol* 31: 606–613.

[Google Scholar](#) | [SAGE Journals](#) | [ISI](#)

Prentice, AM, Jebb, SA (2001) Beyond body mass index. *Obesity Rev* 2: 141–147.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Rucker, JJ, Iliff, J, Nutt, DJ (2018) Psychiatry & the psychedelic drugs. Past, present & future. *Neuropharmacology* 142: 200–218.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

SAMHSA (2019) 2018 National Survey On Drug Use and Health – Public Use File Codebook. Available at: https://www.datafiles.samhsa.gov/sites/default/files/field-uploads-protected/studies/NSD_UH-2018/NSDUH-2018-datasets/NSDUH-2018-DS0001/NSDUH-2018-DS0001-info/NSDUH-2018-D_S0001-info-codebook.pdf

[Google Scholar](#)

Sessa, B (2018) The 21st century psychedelic renaissance: Heroic steps forward on the back of an elephant. *Psychopharmacology* 235: 551–560.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Sexton, JD, Nichols, CD, Hendricks, PS (2019a) Population survey data informing the therapeutic potential of classic and novel phenethylamine, tryptamine, and lysergamide psychedelics. *Front Psychiatry* 10: 896.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Sexton, JD, Crawford, MS, Sweat, NW, et al. (2019b) Prevalence and epidemiological associates of novel psychedelic use in the United States adult population. *J Psychopharmacol* 33: 1058–1067.

[Google Scholar](#) | [SAGE Journals](#) | [ISI](#)

Szabo, A (2015) Psychedelics and immunomodulation: Novel approaches and therapeutic opportunities. *Front Immunol* 6: 358.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Szabo, A (2019) Effects of psychedelics on inflammation and immunity. In: Michael, JW, Ben, S (eds) *Advances in Psychedelic Medicine: State-of-the-Art Therapeutic Applications*. Santa Barbara, CA: ABC-CLIO, pp. 193–213.

[Google Scholar](#)

Szabo, A, Kovacs, A, Frecska, E, et al. (2014) Psychedelic N,N-dimethyltryptamine and 5-methoxy-N,N-dimethyltryptamine modulate innate and adaptive inflammatory responses through the sigma-1 receptor of human monocyte-derived dendritic cells. *PLoS One* 9: e106533.

[Google Scholar](#) | [Crossref](#) | [Medline](#) | [ISI](#)

Thompson, C, Szabo, A (2020) Psychedelics as a novel approach to treating autoimmune conditions. *Immunol Lett* 228: 45–54.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Tourino, MC, de Oliveira, EM, Bellé, LP, et al. (2013) Tryptamine and dimethyltryptamine inhibit indoleamine 2,3 dioxygenase and increase the tumor-reactive effect of peripheral blood mononuclear cells. *Cell Biochem Function* 31: 361–364.

[Google Scholar](#) | [Crossref](#) | [Medline](#) | [ISI](#)

Van Amsterdam, J, Opperhuizen, A, Koeter, M, et al. (2010) Ranking the harm of alcohol, tobacco and illicit drugs for the individual and the population. *Eur Addiction Res* 16: 202–207.

[Google Scholar](#) | [Crossref](#) | [Medline](#) | [ISI](#)

Watts, R, Day, C, Krzanowski, J, et al. (2017) Patients’ accounts of increased “connectedness” and “acceptance” after psilocybin for treatment-resistant depression. *J Humanistic Psychol* 57: 520–564.

[Google Scholar](#) | [SAGE Journals](#) | [ISI](#)

Winkelman, M (2014) Psychedelics as medicines for substance abuse rehabilitation: Evaluating treatments with LSD, Peyote, Ibogaine and Ayahuasca. *Curr Drug Abuse Rev* 7: 101–116.

[Google Scholar](#) | [Crossref](#) | [Medline](#)

Winkelman, MJ, Sessa, B (eds) (2019) *Advances in Psychedelic Medicine: State-of-the-art Therapeutic Applications*. Santa Barbara, CA: ABC-CLIO.

[Google Scholar](#)